

AVIATOR[®] HERBICIDE

SAME ACTIVES AS PULSAR[®] + RUMOUR[®]

Dicamba + Fluroxypyr + Thifensulfuron + Tribenuron Herbicide

WHY USE AVIATOR?

- AgraCity Exclusive Product – Same Actives as Pulsar and Rumour
- Proven control of cleavers, hemp-nettle, redroot pigweed, wild buckwheat
- Controls or suppresses 15 broadleaf weeds
- Use in all soil zones
- Three systemic modes of action for dependable control + resistance management
- Early window of application
- Tank mixable with leading grass herbicides
- No re-cropping restrictions
- Dependable broad-spectrum control of broadleaf weeds in cereal crops at a great price

FOR USE ON

- Spring wheat, durum wheat, barley


WEEDS

CONTROLLED

- Cleavers² (1-4 whorls)
- Corn spurry
- Cow cockle
- Green smartweed
- Hemp-nettle
- Kochia
- Lady's-thumb
- Lamb's quarters
- Redroot pigweed
- Stinkweed
- Tartary buckwheat
- Volunteer canola³
- Wild buckwheat
- Wild mustard

SUPPRESSED

- Canada thistle¹
- Sow thistle (top growth only)



¹ Apply before the bud stage when thistles are <15 cm in height and when majority of thistles have emerged and are actively growing.

² Including Group 2 resistant biotypes.

³ Not including Clearfield[®] tolerant varieties.



AVIATOR HERBICIDE

SAME ACTIVES AS PULSAR + RUMOUR

REQUIRED SURFACTANT

- Agral® 90 or AgSurf® at 2 L per 1,000 L water

ACTIVE INGREDIENT

- Dicamba 480 g/L solution (SN)
- Fluroxypyr 180 g/L emulsifiable concentrate (EC)
- Thifensulfuron methyl 25% + Tribenuron methyl 25% soluble granule (SG)

PACKAGE SIZE

- **Case:** 40 acres

WHEN TO APPLY

- **Crop stage:** 2-5 leaf
- **Weed stage:** 2-3 leaf stage and rosettes 5 cm across

WATER VOLUME

- 10 US gallons/acre (40 L/acre)

RAINFAST PERIOD

- 1 hour

RE-CROPPING OPTIONS

- Grow wheat, barley, oats, rye, canola, flax, forage grasses, lentils, mustard, peas, or fields can be summerfallowed the year after application

TANK MIX OPTIONS*

- **Aurora**®†, **Traxos**®†

STORAGE

- Store in secure dry heated storage. If product is frozen bring to room temperature and agitate before use.

GRAZING AND HARVEST

- Do not graze treated crop or feed to livestock within 7 days of application
- Harvest mature crop 60 days after application

MIXING INSTRUCTIONS

1. Fill the sprayer 1/2 full of clean water, begin agitation and continue throughout mixing and application process.
2. Add the appropriate rate of **Rumour** and dissolve.
3. If required, add the correct amount of grass weed control tank mix partner.
4. Add the required amount of **Ammo**®, followed by **Foxy**®.
5. Add any other required liquid tank mix partner.
6. Add surfactant, then complete filling tank and apply.
7. Product must be applied within 12 hours in order to remain effective.
8. If application is stopped prior to completion, re-suspend spray solution by full agitation prior to commencing spraying again.

UNDERSTANDING PERFORMANCE

- Use 50 mesh or larger screens in spraying equipment.
- Optimal control occurs when applied between 12°C and 24°C in warm moist growing conditions.
- Typical symptoms (discoloration) of dying weeds may not be noticeable for 1-3 weeks after application depending upon growing conditions. Control of weeds may take 3-4 weeks.
- Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.
- Improper or untimely application can result in abnormal bending at the nodes of grain stalks, difficulty in head emergence from the sheath, curled awns, malformed kernels, and sterile florets.
- Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and regrowth may occur.
- Application to crops that are stressed by severe weather conditions, frost, low fertility, water-logged soil (soil at or near field capacity), disease or insect damage may result in crop injury. Drought, disease, or insect damage following application may also result in crop injury.
- Product will degrade if left in the sprayer for more than 24 hours after mixing.
- Dicamba may drift and cause injury to sensitive crops. Do not apply when winds are blowing towards sensitive crops, or during conditions conducive of thermal inversions.

Always read and follow all label precautions and directions.

*When applying tank mix combinations, always follow all label requirements for each product used in the tank mix. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed.

†Supported by AgraCity. For a complete list of tank mix options, go to AgraCity.com/Solutions.

Ammo®, Aurora®, Aviator®, Foxy®, and Rumour® are trademarks of AgraCity or its affiliates. All other products are trademarks of their respective holders.