

FOXXY[®] RCK HERBICIDE


AGRACITY EXCLUSIVE PRODUCT
Fluroxypyr + Thifensulfuron + Tribenuron Herbicide

WHY USE FOXXY RCK?

- AgraCity Exclusive Product - contains a higher rate of active ingredient for outstanding control of kochia (including Group 2 resistant), hemp-nettle, lamb's-quarters and wild buckwheat
- Controls or suppresses 15 broadleaf weeds
- Use in all soil zones
- Two systemic modes of action for resistance management
- Wide window of application
- Excellent crop safety at all application stages
- Great tank mix option with **Aurora[®]** or **Himalaya[®]**
- No re-cropping restrictions
- Proven broad-spectrum control of tough broadleaf weeds at an affordable price

FOR USE ON

- Spring wheat, durum wheat, barley



WEEDS

CONTROLLED

- Cleavers¹ (1-4 whorl)
- Cow cockle
- Green smartweed
- Hemp-nettle
- Kochia¹ (up to 8 leaf)
- Lady's-thumb
- Lamb's quarters
- Redroot pigweed
- Stinkweed
- Volunteer canola²
- Volunteer flax (1-12 cm)
- Wild buckwheat
- Wild mustard

SUPPRESSED

- Canada thistle³
- Stork's-bill (1-8 leaf)

¹ Including Group 2 resistant biotypes

² Not including Clearfield[®] canola

³ Top growth control only. For best results, apply before bud stage when thistles are no taller than 15 cm.



FOXXY RCK HERBICIDE

AGRICITY EXCLUSIVE PRODUCT

REQUIRED SURFACTANT

- Not required if mixed with a wild oat herbicide
- If used alone, Agral® 90 at 2 L per 1,000 L water

ACTIVE INGREDIENTS

- Fluroxypyr 180 g/L emulsifiable concentrate (EC)
- Thifensulfuron methyl 25% + tribenuron methyl 25% Soluble granules (SG)

PACKAGE SIZES

- **Case:** 80 acres
- **Bulk:** 960 acres

WHEN TO APPLY

- **Crop stage:** 2 leaf–flag leaf
- **Weed stage:** less than 10 cm in height or diameter

WATER VOLUME

- 5–10 US gallons/acre (20–40 L/acre)

RAINFEST PERIOD

- 4 hours

RE-CROPPING OPTIONS

- Grow all major crops the year after application.

TANK MIX OPTIONS*

- **Ace®† II, Aurora®†, Hellcat®†, Himalaya®†, MCPA 600 Ester** @190 ml/acre†, **Axial®†, Simplicity™†, Traxos®†, Varro®†**

STORAGE

- Do not freeze. If product is frozen, bring to room temperature and agitate before use.

GRAZING AND HARVEST

- Livestock may graze treated areas 7 days after application
- Withdraw meat animals from treated areas at least 3 days before slaughter
- Allow 7 days after application before cutting hay or harvesting forage
- Mature crops may be harvested 60 days after application.

MIXING INSTRUCTIONS

1. Fill the spray tank 1/2 full of water, start sprayer agitation and continue throughout mixing and application process.
2. Add the required amount of **Rumour™**.
3. Add the required amount of grass weed control tank mix partner.
4. Add the required amount of **Foxyy**.
5. If desired add the required amount of MCPA 600 Ester.
6. Add any required surfactants, complete filling of sprayer and apply.
7. If application is stopped prior to completion, re-suspend spray solution by full agitation prior to commencing spraying again.

UNDERSTANDING PERFORMANCE

- Best results will occur with temperatures between 12° C and 24° C in warm moist conditions when weeds are small and actively growing.
- Ensure thorough spray coverage of target weeds and use higher water volumes with dense crop canopy and/or large weeds.
- Typical symptoms (discoloration, twisting of stems) of dying weeds may not be visible for several days to 2 weeks after application, depending on growing conditions. Control of weeds may take 3–4 weeks.
- Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.
- When weeds are stressed due to drought, flooding or prolonged hot or cool temperatures (less than 8° C or greater than 27° C), control can be reduced or delayed since weeds are not actively growing. Do not make applications to weeds stressed longer than 20 days due to lack of moisture as unsatisfactory control can result.
- Application to crops that are stressed by severe weather conditions, frost, low fertility, water-logged soil (soil at or near field capacity), disease or insect damage may result in crop injury. Drought, disease, or insect damage following application may also result in crop injury.
- Product will degrade if left in the sprayer for more than 24 hours after mixing.

Always read and follow all label precautions and directions.

*When applying tank mix combinations, always follow all label requirements for each product used in the tank mix. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed.

†Supported by AgraCity. For a complete list of tank mix options, go to AgraCity.com/Solutions.

Ace®, Aurora®, Foxyy®, Himalaya®, Hellcat® and Rumour® are trademarks of AgraCity or its affiliates. All other products are trademarks of their respective holders.