

GROUP
2,4

AgraCity

PERFORMANCE
100%
GUARANTEE

FOXXY THIEF™

WHY USE FOXXY THIEF?

- Same actives as Travallas
- Proven control of cleavers, hemp-nettle, redroot pigweed, and lamb's quarters
- Control of group 2 resistant cleavers
- Controls or suppresses 15 broadleaf weeds
- Use in all soil zones
- Systemic dual modes of action for resistance management
- Widest window of application available
- Excellent crop safety at all application stages
- Great tank mix option with **Aurora®, HellCat® or Himalaya™**
- Limited re-cropping restrictions
- Proven broad-spectrum control of the toughest broadleaf weeds in the black soil zone

USE ON

- Spring wheat, durum wheat, barley

WEEDS CONTROLLED

- Chickweed¹
- Cleavers (1-4 whorl)
- Corn spurry
- Cow cockle
- Green smartweed
- Hemp nettle
- Lady's thumb
- Lamb's quarters
- Redroot pigweed
- Russian thistle
- Stinkweed
- Volunteer canola²
- Wild buckwheat³
- Wild mustard

WEEDS SUPPRESSED

- Kochia

¹Apply when the chickweed is small (1 to 6 leaf) and actively growing but before crop canopy prevents thorough coverage of weeds.

²Not including Clearfield varieties

³Apply to actively growing wild buckwheat in the 1 - 3 leaf stage. Under dry conditions control may be reduced. Large plants may regrow after treatment.



CANADIAN
OWNED

SAME ACTIVE AS TRAVALLAS®

Fluroxypyr + Thifensulfuron methyl + Metsulfuron-methyl Herbicide

REQUIRED SURFACTANT

- Not required if mixed with a wild oat herbicide
- If used alone, Icon™, Agra® 90 or AgSurf at 2 L per 1000 L water

ACTIVE INGREDIENTS

- Fluroxypyr 180 g/L emulsifiable concentrate (EC)
- Thifensulfuron-methyl 75% dry flowable (DF)
- Metsulfuron-methyl 60% water dispersible granules (WDG)

PACKAGE SIZES

- **Case:** 80 acres

WHEN TO APPLY

- **Crop stage:** 2 leaf – flag leaf
- **Weed stage:** less than 10 cm in height or diameter unless otherwise noted.

WATER VOLUME

- 10 US gallons/acre (40 L/acre)

RAINFAST PERIOD

- 4 hours

RE-CROPPING OPTIONS

- **10 months after application:** barley, canola, durum wheat, fababean, field corn, flax, oats, peas, spring wheat, soybeans.
- **22 months after application:** above crops plus lentils

SUPPORTED TANK MIXES*

- **Aurora, Hellcat, Himalaya™, MPower® MCPA Ester @190 mL/acre, Axial®, Simplicity™, Traxos®**

STORAGE

- Do not freeze. If product is frozen, bring to room temperature and agitate before use.

GRAZING AND HARVEST

- Livestock may graze treated areas 7 days after application
- Withdraw meat animals from treated areas at least 3 days before slaughter
- Allow 7 days after application before cutting hay or harvesting forage
- Pre-harvest interval: 60 days

MIXING INSTRUCTIONS

1. Fill the spray tank 1/2 full of water, start sprayer agitation and continue throughout mixing and application process.
2. Add the required amount of **Rambo™**.
3. Add the required amount of **MPower® Pro**
4. Add the required amount of grass weed control tank mix partner.
5. Add the required amount of **Foxy™**.
6. If desired add the required amount of **MPower MCPA Ester**.
7. Add any required surfactants, complete filling of sprayer and apply.
8. If application is stopped prior to completion, re-suspend spray solution by full agitation prior to commencing spraying again.

UNDERSTANDING PERFORMANCE

- Best results will occur with temperatures between 12° C and 24° C in warm moist conditions when weeds are small and actively growing.
- Ensure thorough spray coverage of target weeds and use higher water volumes with dense crop canopy and/or large weeds.
- Typical symptoms (discoloration, twisting of stems) of dying weeds may not be visible for several days to 2 weeks after application, depending on growing conditions. Control of weeds may take 3-4 weeks.
- Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.
- When weeds are stressed due to drought, flooding or prolonged hot or cool temperatures (less than 8 °C or more than 27° C), control can be reduced or delayed since weeds are not actively growing. Do not make applications to weeds stressed longer than 20 days due to lack of moisture as unsatisfactory control can result.
- Application to crops that are stressed by severe weather conditions, frost, low fertility, water-logged soil (soil at or near field capacity), disease or insect damage may result in crop injury. Drought, disease, or insect damage following application may also result in crop injury.
- Product will degrade if left in the sprayer for more than 24 hours after mixing.

Always read and observe all label directions including rates, restrictions, and grazing and harvest limitations.

*Supported by AgraCity under PMRA tank mix guidelines. When applying tank mix combinations, read and observe all label directions for each product used in the tank mix, and follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.