

FUSARO™ FUNGICIDE

GROUP
3


SAME ACTIVES AS PROSARO®
Prothioconazole + Tebuconazole Fungicide

WHY USE FUSARO?

- Excellent protection from fusarium head blight
- Tank-mixable with foliar products
- Multi-mode of action
- Excellent value and price with no programming
- Two different systemic active ingredients
- Broad spectrum leaf disease control
- Preventative and curative properties
- Protects grade and quality while increasing yield

FOR USE ON

- Spring wheat, durum wheat, winter wheat and barley



DISEASES CONTROLLED

Spring wheat, durum and winter wheat

- Rusts (leaf, stem and stripe)
- Septoria leaf and glume blotches
- Tab spot
- Powdery mildew

Barley

- Rusts (leaf, stem and Stripe)
- Septoria (net, leaf and spot blotch)
- Scald
- Powdery mildew

DISEASES SUPPRESSED

Spring wheat, durum and winter wheat

- Fusarium head blight



FUSARO FUNGICIDE

SAME ACTIVE AS PROSARO

REQUIRED SURFACTANT

- None required

ACTIVE INGREDIENTS

- Prothioconazole 125 g/L (EC)
- Tebuconazole 125 g/L (EC)

PACKAGE SIZE

- **Case:** 2 x 13 litre
- **Bulk:** 104 litre drum and 625 litre tote

APPLICATION TIMING (CROP)

- **Wheat: spring, durum, winter**
Fusarium head blight: within the time-period when 75% of wheat heads are fully emerged to when 50% of heads on the main stems are in flower. This timing will also control the listed leaf diseases
- **Barley**
Fusarium head blight apply between 70% and 100% full head emergence on the main stem up until 3 days after. This timing will also control the listed leaf diseases

WATER VOLUME

- **Ground:** Minimum 10 US gallons/acre (40 L/acre)
- **Aerial:** Minimum 5 US gallons/acre (20L/acre)

RAINFAST PERIOD

- 1 hour

RE-CROPPING OPTIONS

- Grow all major crops the year after application.

TANK MIX OPTIONS

- None

STORAGE

- Do not freeze. Store in a cool, dry, locked, well-ventilated area without a floor drain. Shake well before use.

GRAZING AND HARVEST

- Allow 5 days after application before cutting hay or harvesting forage.
- Mature crop may be harvested 36 days after application.

MIXING INSTRUCTIONS

1. Fill the spray tank 1/2 full of water and begin agitation.
2. Add the required quantity of **Fusaro** fungicide to the water and complete filling with water to the required total volume.
3. Maintain agitation throughout mixing and spraying.
4. If application is stopped prior to completion, re-suspend spray solution by full agitation prior to commencing spraying again.
5. Clean out sprayer equipment thoroughly after application is complete.

UNDERSTANDING PERFORMANCE

- Consult the product label for application rates and water volumes for all crops and diseases.
- Fusarium head blight risk is greater when the weather is warm and wet at the flowering to soft dough stages. Consult the provincial disease risk maps.
- **Fusaro** fungicide has preventative and curative properties but works best if applied at the onset of disease.
- Good coverage is essential for effective disease control.
- For fusarium head blight, ensure thorough coverage of all wheat heads for optimal suppression. Improved spray coverage of heads may be achieved by using nozzles designed to spray in two directions.
- Do not apply **Fusaro** more than once per crop season. Avoid applying multiple fungicides within the same group during the same crop year to delay the development of resistance.

Always read and follow all label precautions and directions.

*When applying tank mix combinations, always follow all label requirements for each product used in the tank mix. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed.

[†]Supported by AgraCity.

Fusaro® is a trademarks of AgraCity or its affiliates. All other products are trademarks of their respective holders.