

# REVENGE™ B

## WHY USE REVENGE B?

- Same active ingredients as Conquer
- Proven pre-seed control of Lamb's-quarters and Redroot pigweed
- Controls or suppresses 32 annual grass and broadleaf weeds when mixed with glyphosate
- Faster burndown than bromoxynil alone
- Use in all soil zones
- 2 contact modes of action
- No re-cropping restrictions
- Broad-spectrum pre-seed control prior to seeding canola at the best price

## USE ON

- Prior to seeding canola

## WEEDS CONTROLLED

(when mixed with glyphosate acid equivalent at 0.5 – 1.0 L/acre (360 gm a.i./L formulation))

### Grass Weeds: Up To 8 cm In Height Unless Otherwise Noted

- Annual blue grass <sup>3,4</sup>
- Crab grass <sup>3,4</sup>
- Downy brome <sup>2,4</sup>
- Green foxtail
- Giant foxtail <sup>1,4</sup>
- Persian darnel <sup>2,4</sup>
- Volunteer barley <sup>1</sup>
- Volunteer wheat <sup>1</sup>
- Wild oats <sup>1</sup>

### Broadleaf Weeds: Up To 8 cm In Height Unless Otherwise Noted

- Annual sow thistle <sup>3,4</sup>
- Canada fleabane <sup>2</sup>
- Cleavers <sup>2</sup>
- Common ragweed <sup>2</sup>
- Eastern nightshade <sup>7</sup>
- Flixweed <sup>2,4</sup>
- Hemp-nettle <sup>2,4</sup>
- Kochia <sup>3</sup>
- Lady's thumb <sup>1</sup>
- Lamb's-quarters <sup>5</sup>
- Morning glory <sup>6</sup>
- Narrow-leaved hawk's beard <sup>2</sup>
- Prickly lettuce <sup>3,4</sup>
- Redroot pigweed <sup>6</sup>
- Russian thistle <sup>2,4</sup>
- Shepherd's purse <sup>3,4</sup>
- Stinkweed <sup>1</sup>
- Tall waterhemp <sup>7</sup>
- Volunteer canola <sup>1,8</sup> (not including glyphosate tolerant)
- Velvetleaf
- Volunteer flax <sup>2,4</sup>
- Wild buckwheat <sup>2,8</sup>
- Wild mustard <sup>1</sup>

<sup>1</sup>Controlled with 180 g ae glyphosate

<sup>2</sup>Controlled with 180-275 g ae glyphosate

<sup>3</sup>Controlled with 325 g ae glyphosate

<sup>4</sup>Up to 15 cm tall

<sup>5</sup>Up to 7.5 cm tall

<sup>6</sup>Up to 3 leaf

<sup>7</sup>Up to 5 cm tall

<sup>8</sup>1 to 4 leaf



# SAME ACTIVE AS CONQUER® Carfentrazone + Bromoxynil Herbicide

## REQUIRED SURFACTANT

- None required when used alone

## ACTIVE INGREDIENT

- Carfentrazone 240 g/L emulsifiable concentrate (EC)
- Bromoxynil 280 g/L EC

## PACKAGE SIZE

- **Case:** 80 acres

## WHEN TO APPLY

- Immediately prior to seeding, prior to the 4 leaf stage

## WATER VOLUME

- 10 US gallons/acre (40 L/acre)

## RAINFAST PERIOD

- 2 hours

## RE-CROPPING OPTIONS

- Grow all major crops the year after application

## SUPPORTED TANK MIXES\*

- **Disruptor® 360 and 540 glyphosate**

## STORAGE

- If frozen, warm to room temperature and agitate before use

## GRAZING AND HARVEST

- Do not graze the treated crop or cut for feed
- Harvest mature crop 3 days after application

## MIXING INSTRUCTIONS

1. Fill the sprayer 1/2 full of clean water, begin agitation and continue throughout mixing and application process.
2. If required add the correct amount of any dry or solution concentrate products
3. Add the required amount of **Revenge**.
4. Add the required amount of **Buck®**
5. Add the required amount of **Disruptor** glyphosate.
6. Add surfactant if required, then complete filling tank and apply.
7. If application is stopped prior to completion, re-suspend spray solution by full agitation prior to commencing spraying again.

## UNDERSTANDING PERFORMANCE

- Optimal control occurs when applied in warm moist growing conditions when weeds are small and actively growing.
- **Revenge B** is a contact herbicide combination and must be applied directly on growing weeds.
- For best results, spray in the heat of the day with adequate water volume for good coverage, and use a combination of nozzle type and sprayer speed for medium droplet size, adjust sprayer speed to optimize weed coverage
- Typical symptoms of control (browning) appear rapidly after application. Susceptible plants become malformed, with complete control occurring within a few days.
- Because **Revenge B** has no residual, weeds that emerge after application will not be controlled.
- Weed control may be reduced if weeds are stressed by conditions such as drought, waterlogged soils, or excessively cold or hot temperatures.

Always read and observe all label directions including rates, restrictions, and grazing and harvest limitations.

\*Supported by AgraCity under PMRA tank mix guidelines. When applying tank mix combinations, read and observe all label directions for each product used in the tank mix, and follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.