

# TORNADO<sup>®</sup> PRO FUNGICIDE

AGRACITY EXCLUSIVE PRODUCT  
Tebuconazole + Pyraclostrobin Fungicide

## WHY USE TORNADO PRO?

- AgraCity Exclusive Product
- Multiple modes of action for resistance management
- Preventative and curative properties
- Helps protect the crop to increase yield potential
- Improves overall plant health
- Use in all soil zones
- Wide window of application
- Tank mix with foliar
- Proven cereal leaf disease control at an affordable price

## FOR USE ON

- Wheat, barley

## DISEASES

### CONTROLLED

**Spring wheat, winter wheat, durum wheat**

- Rusts (leaf, stem, stripe)
- Septoria leaf blotch
- Tan spot

#### Barley

- Net blotch
- Spot blotch
- Scald
- Rusts (leaf, stem, stripe)
- Septoria leaf blotch
- Powdery mildew



# TORNADO PRO FUNGICIDE

AGRICITY EXCLUSIVE PRODUCT

## REQUIRED SURFACTANT

- None required

## ACTIVE INGREDIENT

- Tebuconazole 250 g/L emulsion in water (EW)
- Pyraclostrobin 250 g/L emulsifiable concentrate (EC)

## PACKAGE SIZE

- **Case:** 9.8 L Tornado + 7.7 L Spade
- **Bulk:** 118 L Tornado + 92.4 L Spade

## APPLICATION TIMING

- Apply at the first sign or very early stage of disease, especially if weather conditions are conducive to disease development

## WATER VOLUME

- **Ground:** Minimum 10 US gallons/acre (40 L/acre)
- **Aerial:** Minimum 5 US gallons/acre (20 L/acre)

## RAINFAST PERIOD

- 1 hour

## RE-CROPPING OPTIONS

- Grow all major crops the year after application

## TANK MIX OPTIONS\*

- None

## STORAGE

- Do not freeze. Store in a cool, dry, well-ventilated place without a floor drain and make sure to avoid cross-contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

## GRAZING AND HARVEST

- Graze or feed treated crops 6 days after application

## PRE-HARVEST INTERVAL

(see label for complete listing)

- **Wheat, barley:** 36 days after application

## MIXING INSTRUCTIONS

1. Fill the spray tank 1/2 full of water, start sprayer agitation, and continue agitation throughout mixing and spraying process.
2. Add required amount of **Tornado** to the tank, followed by **Spade**<sup>®</sup>.
3. Finish filling the tank with water and apply product.
4. If application is stopped prior to completion, re-suspend spray solution by full agitation prior to commencing spraying again.
5. Clean out sprayer equipment thoroughly after application is complete.

## UNDERSTANDING PERFORMANCE

- Consult the product label for uses, application rates and water volumes for all crops and diseases.
- **Tornado Pro** fungicide has preventative and curative properties, but works best if applied preventatively at the onset of disease
- Good spray coverage is critical for effective disease control. Use higher water volumes with dense crop canopies
- Apply **Tornado Pro** only once per crop season. If disease persists or weather conditions are favourable to disease, apply another fungicide product 10-14 days later. Avoid applying products containing the same fungicide groups as **Tornado Pro** to reduce the risk of resistant forms of disease.
- For best disease control, use **Tornado Pro** as part of an integrated disease management approach including resistant seed varieties, crop rotation, and other such management practices.
- Do not apply **Tornado Pro** when the crop is flowering. Applications of fungicides from the strobi class are thought to increase the risk of fusarium head blight if applied during flowering.

Always read and follow all label precautions and directions.

\*When applying tank mix combinations, always follow all label requirements for each product used in the tank mix. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed.

Spade<sup>®</sup> and Tornado<sup>®</sup> are trademarks of AgraCity or its affiliates. All other products are trademarks of their respective holders.